



# Air Impingement Drying of Macroalgae *Sargassum muticum*: effects of the Temperature and the Air-Drying Velocity on the Drying Kinetics

Jeanne Le Loeuff, V. Boy, Pascal Morancais, Nathalie Bourgougnon,  
Jean-Louis Lanoiselle

## ► To cite this version:

Jeanne Le Loeuff, V. Boy, Pascal Morancais, Nathalie Bourgougnon, Jean-Louis Lanoiselle. Air Impingement Drying of Macroalgae *Sargassum muticum*: effects of the Temperature and the Air-Drying Velocity on the Drying Kinetics. International Congress of Chemical and Process Engineering 2021, Mar 2021, Virtual conference, France. hal-04764679

HAL Id: hal-04764679

<https://ubs.hal.science/hal-04764679v1>

Submitted on 4 Nov 2024

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/357535137>

# Air Impingement Drying of Macroalgae *Sargassum muticum*: effects of the Temperature and the Air-Drying Velocity on the Drying Kinetics

Conference Paper · March 2021

CITATIONS

0

READS

46

5 authors, including:



Jeanne Le Loeuff

Upcyclink

16 PUBLICATIONS 13 CITATIONS

[SEE PROFILE](#)



Virginie Boy

Université Bretagne Sud

49 PUBLICATIONS 125 CITATIONS

[SEE PROFILE](#)



Nathalie Bourgougnon

Université Bretagne Sud

161 PUBLICATIONS 5,160 CITATIONS

[SEE PROFILE](#)



Jean-Louis Lanoisellé

Université Bretagne Sud

162 PUBLICATIONS 3,192 CITATIONS

[SEE PROFILE](#)

# Air Impingement Drying of Macroalgae *Sargassum muticum*: effects of the Temperature and the Air-Drying Velocity on the Drying Kinetics

J. Le Loeuff<sup>1</sup>, V. Boy<sup>1</sup>, P. Morançais<sup>2</sup>, N. Bourgougnon<sup>2</sup>, J.-L. Lanoiselle<sup>1</sup>

<sup>1</sup>Univ. Bretagne Sud, IRDL, UMR CNRS 6027, 56300 Pontivy, France; tel. +33 297276770,  
e-mail: jeanne.le-loeuff@univ-ubs.fr

<sup>1</sup>Univ. Bretagne Sud, IRDL, UMR CNRS 6027, 56300 Pontivy, France; tel. +33 297276773,  
e-mail: virginie.boy@univ-ubs.fr

<sup>2</sup>Univ. Bretagne Sud, LBCM, EA 3884, 56300 Pontivy, France; tel. +33 297274993, e-mail:  
pascal.morancais@univ-ubs.fr

<sup>2</sup>Univ. Bretagne Sud, LBCM, EA 3884, 56017 Vannes, France; tel. +33 297017155, e-mail:  
nathalie.bourgougnon@univ-ubs.fr

<sup>1</sup>Univ. Bretagne Sud, IRDL, UMR CNRS 6027, 56300 Pontivy, France; tel. +33 297276774,  
e-mail: jean-louis.lanoiselle@univ-ubs.fr

## 1. Introduction

The brown alga *Sargassum*, which is an invasive species, causes public health troubles because of their massive groundings on beaches. About 10 000 wet tonnes of seaweeds have been grounded on beaches of Caribbean islands daily<sup>1</sup>. One of the main issues to overcome the recovery of this raw material is the seasonal nature of groundings. In this context, the goal of this work is to add value to this alga biomass through the development of a biorefinery to use the resource for longer periods. Owing to their high moisture content, this raw material should be stabilized before being used as biorefinery input. The drying aims to obtain a stabilized product available the entire year. Thermal drying is a complex process and energy intensive. Therefore, modern drying technology focuses on energy saving and on the quality of the product. Air impingement technology aims to enhance the hot air-drying process. Thanks to this technology, the air impingement dryer permits to decrease the drying time and the temperature levels<sup>2</sup>.

## 2. Methods

A prototype of air impingement dryer designed by CIMS company (Sablé-sur-Sarthe, France) was used. The alga *Sargassum muticum* was dried in different conditions. The experiments (quadruplicate) were carried out at four moderate temperatures (40, 50, 60 and 70 °C) at 35 m s<sup>-1</sup> and at the constant temperature of 40 °C at five air-drying velocities (7, 14, 21, 28 and 35 m s<sup>-1</sup>).

### 3. Results and discussion




Fig. 1.  $\ln D_{\text{eff}}$  vs air velocity at  $40^\circ\text{C}$  and temperature at  $35 \text{ m s}^{-1}$  (marks: data; solid line: Arrhenius model for temperature; dotted line: eyes guideline for air velocity)

### 4. Conclusion

This study shows that the apparent moisture diffusivity decreases with the drying temperature and depends on the air velocity (Fig. 1). The lower value of  $D_{\text{eff}}$  is found at  $21 \text{ m s}^{-1}$ , which may be attributed to a crusting phenomenon at high air velocity. This hypothesis is under investigation through rehydration kinetics of the dried algae.

### References

- [1] J.J. Milledge, P.J. Harvey, Golden Tides: Problem or golden opportunity? The valorization of *Sargassum* from beach inundations, Journal of Marine Science and Engineering 2016, 4, 60.
- [2] V. Boy, W. Ben Khalifa, L. Drévillon, Y. Lemée, T. Lendormi, J.-L. Lanoiselé, Air impingement and intermittent drying: application to apple and to mango, Canadian Journal of Chemical Engineering 99 (1), 120-134, 2021.

To reach a moisture content of 10 % (wet basis), the drying time, which decreases with the temperature, varied between 25 and 130 min.

The Fick's diffusional model was fitted to the drying kinetics and the model was applied to determine the apparent moisture diffusivity,  $D_{\text{eff}}$ . For all the conditions,  $D_{\text{eff}}$  varies between  $3.09 \cdot 10^{-10}$  and  $5.14 \cdot 10^{-10} \text{ m s}^{-1}$ . Ten semi-empirical models were applied on the experimental data. The quality of the models was evaluated by four statistical parameters: the adjusted coefficient of determination ( $R^2$ ), the residual sum of squares (RSS), the chi square ( $\chi^2$ ) and the root-mean-square-error (RMSE). At  $40^\circ\text{C}$  and  $35 \text{ m s}^{-1}$ , the Henderson Pabis modified model best fits the experimental data ( $R^2 > 0.99$ ;  $\chi^2 = 5.32 \cdot 10^{-4}$ ;  $\text{RSS} = 6.36 \cdot 10^{-3}$ ;  $\text{RMSE} = 1.71 \cdot 10^{-2}$ ).